

S-ARITHMETIC GROUPS OF SL_2 TYPE

R. SARMA

ABSTRACT. For a number field K , we show that any S -arithmetic subgroup of $SL_2(K)$ contains a subgroup of finite index generated by three elements if $\text{card}(S) \geq 2$.

1. INTRODUCTION AND NOTATION

Let K be a number field and let S_∞ be the set of all nonconjugate embeddings of K into $\mathbb{C}$. We refer to these embeddings as *infinite primes* of K . If r_1 (resp. r_2) is the number of distinct real (resp. nonconjugate complex) embeddings so that the cardinality of S_∞ is $r_1 + r_2$, then $r_1 + 2r_2 = [K : \mathbb{Q}]$, the extension degree of K . Let S be a finite set of primes in K containing S_∞ . The ring of integers in K is denoted by $\mathcal{O}_K$. For a prime ideal $\mathfrak{p}$ of $\mathcal{O}_K$, denote by $v_{\mathfrak{p}}$ the valuation defined by $\mathfrak{p}$. The ring $\mathcal{O}_S := \{x \in K : v_{\mathfrak{p}}(x) \geq 0 \text{ for every prime } \mathfrak{p} \notin S\}$ is called the ring of S -integers of K . Then $\mathcal{O}_{S_\infty} = \mathcal{O}_K$. If F is a subfield of K , then set

$$(1) \quad S(F) := \{\mathfrak{p} \cap \mathcal{O}_F : \mathfrak{p} \in S - S_\infty\} \sqcup S_\infty(F)$$

where $S_\infty(F)$ denotes the infinite primes of F . We write

$$(2) \quad \mathcal{O}_{S(F)} := \{x \in F : v_{\mathfrak{p}}(x) \geq 0 \ \forall \mathfrak{p} \notin S(F)\}$$

the ring of $S(F)$ -integers in F .

For two subgroups H_1 and H_2 in a group, if $H_1 \cap H_2$ is a subgroup of finite index both in H_1 and H_2 , then we say that H_1 and H_2 are *commensurable* and we write $H_1 \asymp H_2$. In particular, a group is commensurable with its subgroups of finite index. A subgroup Γ of $SL_2(K)$ is called *S -arithmetic* if $\Gamma \asymp SL_2(\mathcal{O}_S)$.

A subset X of a group G is called a set of *virtual generators* of G if the group generated by X is a subgroup of finite index in G and the group G is said to be generated *virtually* by X .

Let the cardinality of any set X be denoted by $\text{card}(X)$.

1991 *Mathematics Subject Classification*. Primary 20F05, 11F06; Secondary 22E40

R. Sarma. Harish-Chandra Research Institute, Chhatnag Road, Jhansi, Allahabad 211 019, India. ritumoni@mri.ernet.in.

A number field is called a *totally real* field if all its embeddings are real. A number field is called a *CM field* if it is an imaginary quadratic extension of a totally real field. If a number field is not CM then we refer to it as a *non-CM* field.

For any commutative ring A , denote by

$$(3) \quad \begin{pmatrix} 1 & A \\ 0 & 1 \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} 1 & 0 \\ A & 1 \end{pmatrix})$$

the subgroup of $SL_2(A)$ consisting of matrices of the form

$$\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}) \quad \text{for } x \in A.$$

Let G be any group and let $a, b \in G$. Denote by ${}^a b$ the element aba^{-1} in G .

We use a few well known number theoretic results (for details, see [1],[2]): The ring $\mathcal{O}_K$ of integers in K is a **Dedekind domain**. An ideal of $\mathcal{O}_K$ has a unique **factorization into prime ideals** of $\mathcal{O}_K$. For a finitely generated abelian group H , let $\text{rank}(H)$ denote the **rank** of H as a $\mathbb{Z}$ -module. **Dirichlet's unit theorem** asserts that

$$(4) \quad \text{rank}(\mathcal{O}_K^*) = r_1 + r_2 - 1$$

where r_1 and r_2 are defined as above. Also (cf. Lemma 5)

$$(5) \quad \text{rank}(\mathcal{O}_S^*) = \text{card}(S) - 1.$$

The group of units of a ring A is denoted by A^* . For an ideal $\mathfrak{a}$ of $\mathcal{O}_K$, let the **order** of the class of $\mathfrak{a}$ in the **ideal class group** of K be denoted by $\text{ord}_K(\mathfrak{a})$. It is well known that the class group of a number field is finite. Thus $\text{ord}_K(\mathfrak{a})$ is always a **finite** number.

Now we state the main result of the paper.

Theorem 1. *Let K be a number field and let S be a finite set of primes in K containing the infinite ones such that $\text{card}(S) \geq 2$. Any S -arithmetic subgroup of $SL_2(K)$ is virtually generated by three elements.*

We postpone the proof of this theorem to section 3. In [3], it is shown that the **higher rank arithmetic groups** are virtually generated by three elements. The tools used to prove this do not seem to work for the case of S -arithmetic groups. For instance, if U is a **unipotent group**, and if Γ is a **Zariski dense** subgroup of an arithmetic subgroup of U , then Γ is also arithmetic. This fact plays a crucial role in the case of higher rank arithmetic groups. The analogous statement does not hold in the case of S -arithmetic subgroups. So it needs a separate treatment. The case of SL_2 is the first case that one would like to deal with because this is the simplest possible case. The technique

here may indicate how to proceed for other S -arithmetic groups. In the next section we prove a number theoretic result which asserts that $\mathcal{O}_S$ is **almost** generated by some unit (in fact, by any positive power of that unit) in $\mathcal{O}_S$. Then our main result follows from a theorem due to Vaserstein. The condition that $\text{card}(S) \geq 2$ is equivalent to say that the group $\mathcal{O}_S^*$ is infinite.

2. A NUMBER THEORETIC RESULT

Theorem 2. *Let K be a non-CM field and let S be a finite set of primes including the infinite ones with $\text{card}(S) \geq 2$. Then there exists $\alpha \in \mathcal{O}_S^*$ such that the ring $\mathbb{Z}[\alpha^n]$ is a subgroup of finite index in the ring $\mathcal{O}_S$ of S -integers for every positive integer n .*

The proof of Theorem 2 is divided into a few lemmata.

Lemma 3. *If K is a non-CM field and if F is a proper subfield of K , then $\mathcal{O}_F^*$ is a subgroup of infinite index in $\mathcal{O}_K^*$.*

Proof. Assume to the contrary that there exists a subfield F of K such that the quotient $\mathcal{O}_K^*/\mathcal{O}_F^*$ is finite. We show that K is a CM field and arrive at a contradiction.

Let $d = [K : F]$ be the degree of extension. Let A (resp. B) be the set of all real (resp. nonconjugate complex) embeddings of F over $\mathbb{Q}$. For an embedding $a \in A$, let $x(a)$ (resp. $y(a)$) be the number of real (resp. nonconjugate complex) extensions of a to K . Then we have

$$(6) \quad x(a) + 2y(a) = d.$$

Similarly, we define $x(b), y(b)$ for $b \in B$. Then $x(b) = 0$ and $y(b) = d$. Since $\mathcal{O}_K^*/\mathcal{O}_F^*$ is finite, $\text{rank}(\mathcal{O}_F^*) = \text{rank}(\mathcal{O}_K^*)$. Hence by (4), we have

$$\begin{aligned} \text{card}(A) + \text{card}(B) - 1 &= \sum_{a \in A} x(a) + \left(\sum_{a \in A} y(a) + \sum_{b \in B} y(b) \right) - 1 \\ &= \sum_{a \in A} \{x(a) + y(a)\} + \sum_{b \in B} y(b) - 1 \\ (7) \quad &\geq \text{card}(A) + d \text{card}(B) - 1, \end{aligned}$$

as $x(a) + y(a) \geq 1$. Now since F is a proper subfield of K , we have $d > 1$. Hence using (7), we see that $\text{card}(B) = 0$. Thus, F is a totally real field. Therefore by inequality (7), we get $\text{card}(A) = \sum_{a \in A} \{x(a) + y(a)\}$.

Hence for each $a \in A$, we have

$$(8) \quad x(a) + y(a) = 1.$$

Now, it follows from (6) that $y(a) = d - 1$. Therefore, since $d - 1 \geq 1$, by (8), we have $x(a) = 0$ and $y(a) = 1$. Thus again by (6), we get $d = 2$ so that the field K is an imaginary quadratic extension over the totally real field F which is a contradiction as desired. $\square$

Lemma 4. *Let $K = \mathbb{Q}(\alpha)$ and let α be integral. Then $\mathbb{Z}[\alpha^{-1}]$ is of finite index in $\mathcal{O}_K[\alpha^{-1}]$.*

Proof. Since α is an integral element, we have $\mathbb{Z}[\alpha] \subset \mathbb{Z}[\alpha^{-1}]$. Let n be the index of $\alpha\mathcal{O}_K$ in $\mathcal{O}_K$. For $0 \leq i \leq (n-1)$, the cosets $\alpha\mathcal{O}_K + i$ are the distinct cosets. Otherwise, $\alpha\mathcal{O}_K + i = \alpha\mathcal{O}_K + j$ for $0 \leq i < j \leq (n-1)$ so that $j - i \in \alpha\mathcal{O}_K$. This implies that n divides $j - i$ which is a contradiction. Therefore, $\mathcal{O}_K$ is the union of these n cosets. Thus, in particular,

$$(9) \quad \mathbb{Z}[\alpha] + \alpha\mathcal{O}_K = \mathcal{O}_K.$$

On the other hand, $\mathbb{Z}[\alpha]$ is of finite index in $\mathcal{O}_K$. By (9), we may assume that the distinct cosets (as an additive subgroup) of $\mathbb{Z}[\alpha]$ in $\mathcal{O}_K$ are $\mathbb{Z}[\alpha] + \alpha x_i$ with $0 \leq i \leq (m-1)$. We claim that the representatives of $\mathcal{O}_K[\alpha^{-1}]/\mathbb{Z}[\alpha^{-1}]$ in $\mathcal{O}_K[\alpha^{-1}]$, are αx_i (not necessarily distinct). Let $y \in \mathcal{O}_K$. Then, by (9), $y = y_1 + \alpha x_{i_1}$ for $y_1 \in \mathbb{Z}[\alpha]$ and $0 \leq i_1 \leq (m-1)$. Thus $\alpha^{-1}y = \alpha^{-1}y_1 + x_{i_1}$. Again, using (9), we have $x_{i_1} = z_1 + \alpha x_{i_2}$ for $z_1 \in \mathbb{Z}[\alpha]$ and $0 \leq i_2 \leq (m-1)$ so that $\alpha^{-1}y = (\alpha^{-1}y_1 + z_1) + \alpha x_{i_2}$. Therefore, $\mathbb{Z}[\alpha^{-1}] + \alpha^{-1}y = \mathbb{Z}[\alpha^{-1}] + \alpha x_{i_2}$. Thus inductively, one can show that $\mathbb{Z}[\alpha^{-1}] + \alpha^{-r}y = \mathbb{Z}[\alpha^{-1}] + \alpha x_i$ for some $0 \leq i \leq (m-1)$. $\square$

Lemma 5. *Let K be a number field and let S be a finite set of primes in K containing S_∞ . Assume that $S - S_\infty = \{\mathfrak{q}_1, \dots, \mathfrak{q}_r\}$, $\text{ord}_K(\mathfrak{q}_i) = a_i$ and that $\mathfrak{q}_i^{a_i}$ is generated by $\beta_i \in \mathcal{O}_K \forall i$. Then $\mathcal{O}_S = \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_r^{-1}]$.*

Proof. Obviously, $\mathcal{O}_S \supset \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_r^{-1}]$. To see the other containment, let $x \in \mathcal{O}_S$. Then $x = yz^{-1}$ for $y, z \in \mathcal{O}_K$ and $v_{\mathfrak{p}}(z) = 0$ for $\mathfrak{p} \notin S$ so that, by prime factorization, $z\mathcal{O}_K = \prod_{i=1}^r \mathfrak{q}_i^{n_i}$ with $n_i \geq 0$. Let

$$m = \prod_{i=1}^r a_i. \text{ Since } \mathfrak{q}_i^{a_i} \text{ is generated by } \beta_i, \text{ we have } z^{-m} = u \prod_{i=1}^r \beta_i^{-n'_i} \text{ for}$$

some $u \in \mathcal{O}_K^*$ and $n'_i \geq 0$ so that $z^{-m} \in \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_r^{-1}]$. Further, $z^{-1} = z^{m-1}z^{-m}$ and $z^{m-1} \in \mathcal{O}_K$. Therefore, $z^{-1} \in \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_r^{-1}]$ and hence $x = yz^{-1} \in \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_r^{-1}]$. $\square$

Now by Lemma 4 and Lemma 5, we have the following lemma.

Lemma 6. *Suppose that R is a subring of finite index in $\mathcal{O}_K$. Then with the notation as in Lemma 5, the ring $R[\beta_1^{-1}, \dots, \beta_r^{-1}]$ is of finite index in $\mathcal{O}_S$.* $\square$

Let $\{S_i : 1 \leq i \leq s\}$ be the set of **all the proper subsets** of S and let $\{K_j : 1 \leq j \leq t\}$ be the set of **all the proper subfields** of K . Define

$$(10) \quad V_i := \mathcal{O}_{S_i}^* \otimes_{\mathbb{Z}} \mathbb{Q} \quad \text{and} \quad W_j := (\mathcal{O}_{S(K_j)}^* \cap \mathcal{O}_S^*) \otimes_{\mathbb{Z}} \mathbb{Q}.$$

Then V_i (resp. W_j) is vector space of dimension $\text{rank}(\mathcal{O}_{S_i}^*)$ (resp. $\text{rank}(\mathcal{O}_{S(K_j)}^*)$) over $\mathbb{Q}$. By Lemma 5, we have $\mathcal{O}_S^* = \mathcal{O}_K^* \times \mathbb{Z}^r$ where $r = \text{card}(S) - \text{card}(S_\infty)$.

Lemma 7. *With the above notation, if K is a non-CM field, there exists $\alpha \in \mathcal{O}_S^* - (\bigcup_{i=1}^s V_i) \cup (\bigcup_{j=1}^t W_j)$ such that $v_{\mathfrak{p}}(\alpha) < 0$ for all $\mathfrak{p} \in S - S_\infty$.*

Proof. For each $1 \leq j \leq s$, we have (see (1) and (2) for definition)

$$\begin{aligned} \text{rank}(\mathcal{O}_{S(K_j)}^*) &= \text{card}(S(K_j)) - 1 \\ &= \{\text{card}(S_\infty(K_j)) - 1\} + \text{card}(S(K_j) - S_\infty(K_j)) \\ (11) \quad &= \text{rank}(\mathcal{O}_{K_j}^*) + \text{card}(S(K_j) - S_\infty(K_j)). \end{aligned}$$

Since K is a non-CM field, by Lemma 3, $\text{rank}(\mathcal{O}_{K_j}^*) < \text{rank}(\mathcal{O}_K^*)$. Moreover, $\text{card}(S(K_j) - S_\infty(K_j)) \leq \text{card}(S - S_\infty)$. Therefore, we get

$$(12) \quad \text{rank}(\mathcal{O}_{S(K_j)}^* \cap \mathcal{O}_S^*) < \text{rank}(\mathcal{O}_S^*).$$

Further, $\text{rank}(\mathcal{O}_{S_i}^*) = \text{card}(S_i) - 1 < \text{rank}(\mathcal{O}_S^*)$. Then by comparing the dimensions, we have $V_i \subsetneq V$ and $W_j \subsetneq V$ where $V := \mathcal{O}_S^* \otimes_{\mathbb{Z}} \mathbb{Q}$ (cf. (10)). Since a finite union of proper subspaces of a vector space over an infinite field is a proper subset of the vector space, we have $V - (\bigcup_{i=1}^s V_i) \cup (\bigcup_{j=1}^t W_j)$ is **nonempty**. Let

$$(13) \quad X := \{x \in \mathcal{O}_S^* : v_{\mathfrak{p}}(x) < 0 \forall \mathfrak{p} \in S - S_\infty\}.$$

Then X is Zariski dense in V . Thus the set

$$Y := X - (\bigcup_{i=1}^s V_i) \cup (\bigcup_{j=1}^t W_j)$$

is also nonempty. If $\alpha \in Y$, then $\alpha^n \in Y$. Thus, $\alpha \in \mathcal{O}_S^*$ can be chosen such that $v_{\mathfrak{p}}(\alpha) < 0$ for each $\mathfrak{p} \in S - S_\infty$. $\square$

Lemma 8. *Let α be chosen as in Lemma 7. Then the ring $\mathbb{Z}[\alpha^n]$ is a subgroup of finite index in $\mathcal{O}_S$ for every positive integer n .*

Proof. We claim $\mathbb{Q}(\alpha) = K$. If not, then let $\mathbb{Q}(\alpha) = L$ such that $L \subsetneq K$. Assume for $\mathfrak{p} \notin S$ and $x \in \mathcal{O}_L$ that $v_{\mathfrak{p} \cap \mathcal{O}_L}(x) \neq 0$ so that $x\mathcal{O}_L \subset \mathfrak{p} \cap \mathcal{O}_L$. Then, $x\mathcal{O}_K \subset (\mathfrak{p} \cap \mathcal{O}_L)\mathcal{O}_K \subset \mathfrak{p}$ so that $v_{\mathfrak{p}}(x) \neq 0$. Thus, equivalently, for $x \in \mathcal{O}_L$, if $v_{\mathfrak{p}}(x) = 0$ for every $\mathfrak{p} \notin S$, we have $v_{\mathfrak{p}}(x) = 0$ for every $\mathfrak{p} \notin S(L)$. Therefore, in particular, $v_{\mathfrak{p}}(\alpha^{-1}) = 0 \forall \mathfrak{p} \notin S(L)$

so that $\alpha \in \mathcal{O}_{S(L)}^* \cap \mathcal{O}_S^*$. This contradicts the choice of α . Hence $\mathbb{Q}(\alpha) = K$.

Since $K = \mathbb{Q}(\alpha)$, we also have $K = \mathbb{Q}(\alpha^{-1})$ and since α^{-1} is integral in K , the ring $\mathbb{Z}[\alpha^{-1}]$ is a subgroup of finite index in $\mathcal{O}_K$. Let $S - S_\infty = \{\mathfrak{p}_i : 1 \leq i \leq l\}$. Consider the prime factorization

$$(14) \quad \alpha^{-1} \mathcal{O}_K = \prod_{i=1}^l \mathfrak{p}_i^{n_i}$$

where $n_i \geq 0$. In fact, by our choice of α , $n_i > 0$ for each i . Let $\text{ord}_K(\mathfrak{p}_i) = r_i$ and let $\mathfrak{p}_i^{r_i} = \beta_i \mathcal{O}_K$ for $\beta_i \in \mathcal{O}_K$. Then, we have

$$(15) \quad \alpha^m = \prod_{i=1}^l \beta_i^{-b_i}$$

(if necessary absorbing a unit with β_1) for some integers $m > 0$ and $b_i > 0$. Since $\beta_i \in \mathcal{O}_K$, it follows by (15) that $\beta_i^{-1} \in \mathcal{O}_K[\alpha]$. Now by Lemma 6, the ring $\mathcal{O}_K[\alpha] = \mathcal{O}_S$. Thus, by Lemma 4, the ring $\mathbb{Z}[\alpha]$ is of finite index in $\mathcal{O}_S$. $\square$

This completes the proof of Theorem 2.

In fact, we have proved more.

Corollary 1. *Let K be any finite extension of $\mathbb{Q}$ and let S be as before. If $\text{rank}(\mathcal{O}_{S(L)}^* \cap \mathcal{O}_S^*) < \text{rank}(\mathcal{O}_S^*)$ for every proper subfield L of K , then there exists $\alpha \in \mathcal{O}_S^*$ such that the ring $\mathbb{Z}[\alpha^n]$ is a subgroup of finite index in $\mathcal{O}_S$ for every $n \geq 1$.* $\square$

The hypothesis of Corollary 1 may hold sometimes even for a CM field. Here we see two examples:

Example. (i) The field $K = \mathbb{Q}(\sqrt{-1})$ is a CM field and $\mathcal{O}_K = \mathbb{Z}[\sqrt{-1}]$. The prime ideal $2\mathbb{Z}$ of $\mathbb{Q}$ is totally ramified in K . In fact, $2\mathcal{O}_K = \mathfrak{p}^2$ where $\mathfrak{p} = \langle 1 + \sqrt{-1} \rangle$. Let $S - S_\infty = \{\mathfrak{p}\}$. For K , the set S_∞ of infinite primes is singleton. Thus $\text{card}(S) = 2$ and hence $\text{rank}(\mathcal{O}_S^*) = 1$. Also, $\mathcal{O}_{S(\mathbb{Q})} = \mathbb{Z}[\frac{1}{2}]$ and so $\text{rank}(\mathcal{O}_{S(\mathbb{Q})}^* \cap \mathcal{O}_S^*) = 1$ (observe that $\mathcal{O}_S = \mathbb{Z}[\sqrt{-1}][\frac{1}{1+\sqrt{-1}}]$ includes $\mathcal{O}_{S(\mathbb{Q})}$). This is an example which does not satisfy the hypothesis corollary 1.

(ii) Next consider the ideal $5\mathbb{Z}$ of $\mathbb{Q}$ which splits completely in K : $5\mathcal{O}_K = \mathfrak{p}_1 \mathfrak{p}_2$ where $\mathfrak{p}_1 = \langle 5, 2 + \sqrt{-1} \rangle$ and $\mathfrak{p}_2 = \langle 5, 2 - \sqrt{-1} \rangle$. Let $S - S_\infty = \{\mathfrak{p}_1, \mathfrak{p}_2\}$. Then $\text{card}(S) = 3$ and hence $\text{rank}(\mathcal{O}_S^*) = 2$. The contraction of the primes of $S - S_\infty$ to $\mathbb{Q}$ are $5\mathbb{Z}$ each. Therefore, $\mathcal{O}_{S(\mathbb{Q})} = \mathbb{Z}[\frac{1}{5}]$ and hence $\text{rank}(\mathcal{O}_{S(\mathbb{Q})}^* \cap \mathcal{O}_S^*) = 1$. This is an example of a set of primes of the CM-field K which satisfies the hypothesis.

We need Corollary 1 to prove the main theorem of the paper.

3. PROOF OF THE MAIN THEOREM

We imitate the proof of the same result for the case of arithmetic subgroups of $SL_2(K)$ (cf. [3]). Here, we state a result due to Vaserstein which we use in the proof of Theorem 1.

Theorem 9 (Vaserstein). [5] *Let K be a number field and let S be a finite set of primes in K including S_∞ such that $\text{card}(S) \geq 2$. Let $\mathfrak{a}$ be a nonzero ideal of $\mathcal{O}_S$. The group generated by $\begin{pmatrix} 1 & \mathfrak{a} \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ \mathfrak{a} & 1 \end{pmatrix}$ is a subgroup of finite index in $SL_2(\mathcal{O}_S)$.*

To prove Theorem 1, it suffices to show that any subgroup of finite index in $SL_2(\mathcal{O}_S)$ is virtually generated by three elements. Let Γ be a subgroup of finite index in $SL_2(\mathcal{O}_S)$. Without loss of generality we assume that it is a normal subgroup. Let its index in $SL_2(\mathcal{O}_S)$ be h .

Case 1: The pair (K, S) is such that for **every proper subfield** L of K , we have

$$(16) \quad \text{rank}(\mathcal{O}_{S(L)}^* \cap \mathcal{O}_S^*) < \text{rank}(\mathcal{O}_S^*).$$

Choose $\alpha \in \mathcal{O}_S^*$ as in **Corollary 1**. Obviously, $\begin{pmatrix} \alpha^h & 0 \\ 0 & \alpha^{-h} \end{pmatrix} \in \Gamma$. Since $\mathbb{Z}[\alpha^h]$ is a subring of finite index in $\mathcal{O}_S$, we replace α^h by α and assume that $\gamma := \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \in \Gamma$. Define, $\psi_1 := \begin{pmatrix} 1 & 0 \\ h & 1 \end{pmatrix} \in \Gamma$ and $\psi_2 := \begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix} \in \Gamma$. Let $\Gamma_0 = \langle \gamma, \psi_1, \psi_2 \rangle$. We claim that Γ_0 is a subgroup of finite index in $SL_2(\mathcal{O}_S)$.

Indeed, $\gamma^{-r} \psi_1^s \gamma^r = \begin{pmatrix} 1 & 0 \\ s\alpha^{2r}h & 1 \end{pmatrix} \in \Gamma_0$ and $\gamma^r \psi_2^s \gamma^{-r} = \begin{pmatrix} 1 & s\alpha^{2r}h \\ 0 & 1 \end{pmatrix} \in \Gamma_0$. One concludes from this that Γ contains $\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ y & 1 \end{pmatrix}$ with $x, y \in h\mathbb{Z}[\alpha^2]$. By Corollary 1, $h\mathbb{Z}[\alpha^2]$ is of finite index in the additive group $\mathcal{O}_S$. If m is the index then the ideal $\mathfrak{a} := m\mathcal{O}_S$ is contained in $h\mathbb{Z}[\alpha^2]$. Now it follows from Theorem 9 that the group Γ_0 is a subgroup of finite index in $SL_2(\mathcal{O}_S)$.

Case 2: The pair (K, S) is such that the inequality (16) **does not hold for some proper subfield** F of K . That is, we have

$$(17) \quad \text{rank}(\mathcal{O}_{S(F)}^* \cap \mathcal{O}_S^*) = \text{rank}(\mathcal{O}_S^*).$$

This implies that $\text{rank}(\mathcal{O}_F^*) = \text{rank}(\mathcal{O}_K^*)$. Thus, by Lemma 3, K is a CM field and in fact $K = F(\sqrt{-d})$ so that F is a totally real field and d a totally positive integer in F . Thus, we have

$$(18) \quad \mathcal{O}_{S(F)}^* \asymp \mathcal{O}_S^*,$$

$$(19) \quad \mathcal{O}_F^* \asymp \mathcal{O}_K^*.$$

We prove a number theoretic lemma here.

Lemma 10. *With the above notation, let (17) hold for a CM field $K = F(\sqrt{-d})$. There exists $\alpha \in \mathcal{O}_{S(F)}^* \cap \mathcal{O}_S^*$ such that the ring $\mathbb{Z}[\alpha^n][\sqrt{-d}]$ is of finite index in $\mathcal{O}_S$ for any integer n .*

Proof. In the case of a quadratic extension, a prime ideal of the base field is either inert or totally ramified or split completely (into two distinct primes). We claim that the set $S(F)$ (cf. (1)), does not contain any finite prime which splits completely in K . To the contrary, if $S(F)$ contains a split prime $\mathfrak{q}$ so that $\mathfrak{q}\mathcal{O}_K = \mathfrak{q}_1\mathfrak{q}_2$, then we have two possibilities, namely, $\mathfrak{q}_1, \mathfrak{q}_2 \in S$ or $\mathfrak{q}_1 \in S$ and $\mathfrak{q}_2 \notin S$. If $\mathfrak{q}_1, \mathfrak{q}_2 \in S$, then $\text{card}(S(F)) < \text{card}(S)$ (since $\mathfrak{q}_1$ and $\mathfrak{q}_2$ are contracted to the same prime $\mathfrak{q}$ in F) and thus (17) does not hold. This is a contradiction. Next, assume that $\mathfrak{q}_1 \in S$ and $\mathfrak{q}_2 \notin S$. Let β (resp. γ_1) be the generator of $\mathfrak{q}^{\text{ord}_F(\mathfrak{q})}$ (resp. $\mathfrak{q}_1^{\text{ord}_K(\mathfrak{q}_1)}$). By (18), we have $\mathcal{O}_S \supset \mathcal{O}_{S(F)}$ so that $\beta \in \mathcal{O}_S$. Again (18) and (19) together imply that $\gamma_1^m \in \mathcal{O}_{S(F)}$ for some $m > 0$ so that $\gamma_1^m = u\beta^b x$ for some $b > 0$ and $u \in \mathcal{O}_K^* \cap \mathcal{O}_F^*$ and $x \in \mathcal{O}_{S(F)}^* \cap \mathcal{O}_S^*$ with $v_{\mathfrak{p}}(x) = 0$ for $\mathfrak{p} \notin S(F)$. Then $v_{\mathfrak{q}_2}(\gamma_1) = 0$ whereas $v_{\mathfrak{q}_2}(u\beta^b x) > 0$. This is a contradiction. Therefore, we have

$$(20) \quad (\mathfrak{q} \cap \mathcal{O}_F)\mathcal{O}_K = \mathfrak{q} \text{ or } \mathfrak{q}^2.$$

Let $\text{ord}_F(\mathfrak{q} \cap \mathcal{O}_F) = a$. Then, by (20), we see that $(\mathfrak{q} \cap \mathcal{O}_F)^a \mathcal{O}_K = ((\mathfrak{q} \cap \mathcal{O}_F)\mathcal{O}_K)^a = \mathfrak{q}^a$ or $\mathfrak{q}^{2a}$ is a principal ideal. Thus, $(\mathfrak{q} \cap \mathcal{O}_F)^a$ and $\mathfrak{q}^b$ (for $b = a$ or $2a$) are generated by the same element $\beta \in \mathcal{O}_F$.

Let $S - S_\infty = \{\mathfrak{p}_1, \dots, \mathfrak{p}_s\}$. Choose $\beta_i \in \mathcal{O}_F$ such that $(\mathfrak{p}_i \cap \mathcal{O}_F)^{\text{ord}_F(\mathfrak{p}_i \cap \mathcal{O}_F)} = \beta_i \mathcal{O}_F$. Then, by Lemma 5, $\mathcal{O}_{S(F)} = \mathcal{O}_F[\beta_1^{-1}, \dots, \beta_m^{-1}]$ and $\mathcal{O}_S = \mathcal{O}_K[\beta_1^{-1}, \dots, \beta_m^{-1}]$. Now, since $\mathcal{O}_F[\sqrt{-d}]$ is of finite index in $\mathcal{O}_K$, by Lemma 6, we have $\mathcal{O}_{S(F)}[\sqrt{-d}] = \mathcal{O}_F[\sqrt{-d}][\beta_1^{-1}, \dots, \beta_m^{-1}]$ is of finite index in $\mathcal{O}_S$. Since F is a non-CM field, by Theorem 2, one can choose $\alpha \in \mathcal{O}_{S(F)}^* \cap \mathcal{O}_S^*$ such that $\mathbb{Z}[\alpha^n]$ is of finite index in $\mathcal{O}_{S(F)}$ for every $n \geq 1$. Then $\mathbb{Z}[\alpha^n][\sqrt{-d}]$ is of finite index in $\mathcal{O}_S$. $\square$

Define γ and ψ_1 as in **case 1**. We modify the definition of ψ_2 by $\psi_2 := \begin{pmatrix} 1 & h\sqrt{-d} \\ 0 & 1 \end{pmatrix} \in \Gamma$. Let $\Gamma_0 := \langle \gamma, \psi_1, \psi_2 \rangle$. We show that Γ_0 is a subgroup of finite index in $SL_2(\mathcal{O}_S)$.

Since F is a non-CM field, by an argument similar to case 1, one shows that there is an ideal $\mathfrak{a}$ of $\mathcal{O}_{S(F)}$ such that

$$(21) \quad \begin{pmatrix} 1 & 0 \\ \mathfrak{a} & 1 \end{pmatrix} \subset \Gamma_0 \quad \text{and} \quad \begin{pmatrix} 1 & \sqrt{-d}\mathfrak{a} \\ 0 & 1 \end{pmatrix} \subset \Gamma_0.$$

Then for $x \in \mathfrak{a}$, using Bruhat decomposition (see [4, 8.3]) of ψ_2 , we have

$$(22) \quad {}^{\psi_2} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} = {}^u \begin{pmatrix} 1 & h^2 dx \\ 0 & 1 \end{pmatrix} \in \Gamma_0 \quad \text{where } u = \begin{pmatrix} 1 & 0 \\ \frac{1}{h\sqrt{-d}} & 1 \end{pmatrix}.$$

Let $\mathfrak{b} = h^2 d \mathfrak{a}$. Then we have

$$(23) \quad {}^u \begin{pmatrix} 1 & \mathfrak{b} \\ 0 & 1 \end{pmatrix} \subset \Gamma_0 \quad \text{and} \quad {}^u \begin{pmatrix} 1 & 0 \\ \mathfrak{b} & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \mathfrak{b} & 1 \end{pmatrix} \subset \Gamma_0.$$

Let Γ_1 be the subgroup of $SL_2(\mathcal{O}_F)$ generated by $\begin{pmatrix} 1 & \mathfrak{b} \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ \mathfrak{b} & 1 \end{pmatrix}$. Then, by (23), we have ${}^u \Gamma_1 \subset \Gamma_0$. By Theorem 9, the index of Γ_1 in $SL_2(\mathcal{O}_F)$ is finite. Thus it follows that there exists an integer N such that

$$(24) \quad \gamma^N \in \Gamma_1 \cap \Gamma_0.$$

Since ${}^u \Gamma_1 \subset \Gamma_0$, we have ${}^u \gamma^N \in \Gamma_0$.

Therefore, ${}^u \gamma^{-N} \gamma^N = \begin{pmatrix} 1 & 0 \\ (\alpha^{2N} - 1) \frac{\sqrt{-d}}{hd} & 1 \end{pmatrix} \in \Gamma_0$. Now by conjugating this element and its powers by negative powers of γ , one shows that

$$(25) \quad \Gamma_0 \supset \begin{pmatrix} 1 & 0 \\ \sqrt{-d}\mathfrak{c} & 1 \end{pmatrix}$$

where $\mathfrak{c} := (\alpha^{2N} - 1)\mathbb{Z}[\alpha^2] \cap \mathfrak{a}$. Now $\mathfrak{c} + \sqrt{-d}\mathfrak{c}$ is a subgroup of finite index in $\mathcal{O}_{S(F)}[\sqrt{-d}]$ and hence in $\mathcal{O}_S$. Therefore, the group $\mathfrak{c} + \sqrt{-d}\mathfrak{c}$ contains a nonzero ideal $\mathfrak{q}$ of $\mathcal{O}_S$. Since $\mathfrak{c} \subset \mathfrak{a}$, by (21) and (25), we have

$$(26) \quad \begin{pmatrix} 1 & 0 \\ \mathfrak{q} & 1 \end{pmatrix} \subset \Gamma_0.$$

Again, for $y \in \mathfrak{a}$, using the Bruhat decomposition of ψ_1 , we have

$$(27) \quad {}^{\psi_1} \begin{pmatrix} 1 & y\sqrt{-d} \\ 0 & 1 \end{pmatrix} = {}^{v\varphi} \begin{pmatrix} 1 & 0 \\ h^2 y d & 1 \end{pmatrix} \in \Gamma_0$$

where $v = \begin{pmatrix} 1 & \frac{1}{h} \\ 0 & 1 \end{pmatrix}$ and $\varphi = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sqrt{-d}} \end{pmatrix}$. Thus we have

$$(28) \quad {}^{v\varphi} \begin{pmatrix} 1 & \mathfrak{b} \\ 0 & 1 \end{pmatrix} \subset \Gamma_0, \quad \text{and} \quad {}^{v\varphi} \begin{pmatrix} 1 & 0 \\ \mathfrak{b} & 1 \end{pmatrix} \subset \Gamma_0.$$

Therefore, ${}^{v\varphi}\Gamma_1 \subset \Gamma_0$ and hence ${}^{v\varphi}\gamma^N \in \Gamma_1 \cap \Gamma_0$. Thus, using (24) we have

$$(29) \quad {}^{v\varphi}\gamma^N \gamma^{-N} = \begin{pmatrix} 1 & (1 - \alpha^{2N})\frac{1}{h} \\ 0 & 1 \end{pmatrix} \in \Gamma_0.$$

Again by conjugating this element and its powers by nonnegative powers of γ , one shows that

$$(30) \quad \begin{pmatrix} 1 & \mathfrak{c} \\ 0 & 1 \end{pmatrix} \subset \Gamma_0.$$

Since $\mathfrak{c} \subset \mathfrak{a}$, by (21) and (30), we have

$$(31) \quad \begin{pmatrix} 1 & \mathfrak{q} \\ 0 & 1 \end{pmatrix} \subset \Gamma_0.$$

It follows from (26) and (31), and by Theorem 9, that the group Γ_0 is a subgroup of finite index in $SL_2(\mathcal{O}_S)$. This completes the proof of Theorem 1. $\square$

REFERENCES

- [1] D. A. Marcus, *Number fields*, Springer Verlag, 1977.
- [2] V. Platonov and A. Rapinchuk, *Algebraic Groups and Number Theory*, Academic Press, INC 1991.
- [3] Ritumoni Sarma, T. N. Venkataramana, *Generators of Arithmetic Groups*, (to appear in *Geometriae Dedicata*).
<http://front.math.ucdavis.edu/math.GR/0409345>.
- [4] T. A. Springer, *Linear Algebraic Groups*, Progress in Math. Vol. 9. Birkhauser, second ed., 1998.
- [5] L. N. Vaserstein, *On the Group SL_2 over Dedekind Rings of Arithmetic type*, Math. USSR; Sbornik, **18** (1972), No.2, p. 321-332.

HARISH-CHANDRA RESEARCH INSTITUTE, JHUNSI, CHHATNAG ROAD, ALLAHABAD, INDIA.

E-mail address: ritumoni@mri.ernet.in